

AD-A038 206

JOHNS HOPKINS UNIV BALTIMORE MD DEPT OF BIOMEDICAL E--ETC F/6 6/5
PREDICTIVE MODEL OF THE AUDITORY PROCESS AS RELATED TO COMMUNIC--ETC(U)
FEB 77 M B SACHS F44620-76-C-0066

UNCLASSIFIED

AFOSR-TR-77-0364

NL

1 OF 1
AD
A038206



END

DATE
FILMED
4-77

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ADA 038206

19 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER AFOSR - TR - 77 - 0364	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) PREDICTIVE MODEL OF THE AUDITORY PROCESS AS RELATED TO COMMUNICATION SYSTEMS.		5. TYPE OF REPORT & PERIOD COVERED Final Scientific Report 1 Jan 76 - 31 Dec 76	
6. AUTHOR(s) Murray B. Sachs		7. CONTRACT OR GRANT NUMBER(s) F44620-76-C-0066	
8. PERFORMING ORGANIZATION NAME AND ADDRESS The Johns Hopkins University School of Medicine, Biomedical Engineering Baltimore, Maryland 21205		9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2312A4	
10. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research (NL) Bolling AFB, DC 20332		11. REPORT DATE 15 Feb 1977	
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 7p. 2312 A4		13. NUMBER OF PAGES 5	
14. SECURITY CLASS. (of this report) Unclassified		15. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)			
ABSTRACT (Continue on reverse side if necessary and identify by block number) Studies involving the coding of complex stimuli in the cochlear nerve revealed that recovery functions for auditory nerve discharge rate following one minute exposures to tones resulted in increased rapidity of recovery as test tone intensity increased. Recovery time-constant is a monotonic decreasing function of test intensity and exposure intensity. Recovery of human detection thresholds were measured following similar exposures and it was found that recovery time-constant is a monotonically decreasing function of			

DDC
RECEIVED
APR 8 1977
RESOLVED
C

AD No.

DDC FILE COPY

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

407253

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. Cont'd.

test-tone intensity but is a non-monotonic function of exposure tone intensity. Measurements of rate versus level functions for auditory nerve fibers for tone stimuli indicated there is a continuum of saturation behaviors related to fiber thresholds and that rate-level functions slopes depend on both frequency of tone and threshold of the fibers. A stimulus-response relation for two-tone stimuli was expanded to consider those where the ratio of component amplitudes were held constant. This case correspondes to a continuous spectrum signal (e.g., speech) whose spectrum is held constant while overall amplitude is varied. The first study of coding in the cochlear nerve was completed and published. It showed that the stimulus transformations performed by the avian cochlea are similar in most respects to those performed by the mammalian cochlea. Results dealing with the coding of stimuli in avian cochlear nuclei showed nucleus magnocellularis presents a homogeneous population of cells, whereas cells in the nucleus angularis are more complex and are of at least three types based on response and inhibitory areas.

ACCESSION for		White Section	<input checked="checked" type="checkbox"/>
NTIS		Both Section	<input type="checkbox"/>
G			
UNCLASSIFIED			
CLASSIFICATION			
BY		DISTRIBUTION/AVAILABILITY CODES	
DATE		AVAIL. 204/OF SPECIAL	
A			

PREDICTIVE MODEL OF THE AUDITORY PROCESS
AS RELATED TO COMMUNICATION SYSTEMS

I. Coding of Complex Stimuli in the Cochlear Nerve

All of the work related to this part of the project has been published. A brief statement of results will be given here.

1. Adaptation in the Cochlear Nerve: Physiology and Psychophysics

A. Recovery functions for auditory nerve discharge rate following one-minute exposures to tones were measured with the following results:

- i. Recovery proceeds more rapidly as test tone intensity increases. Recovery time constant is a monotonic decreasing function of test intensity.
- ii. Recovery time-constant is a monotonically decreasing function of exposure intensity.

B. Recovery of human detection thresholds were measured following similar exposures with the following results:

- i. Recovery time-constant is a monotonically decreasing function of test-tone intensity.
- ii. Recovery time-constant is a non-monotonic function of exposure tone intensity, increasing at low levels and decreasing for higher levels.
- iii. The psychophysical results are predicted by a signal detection model based on the physiological results.

References:

- Young, E. and Sachs, M. B., Recovery from sound exposure in auditory-nerve fibers, J. Acoust. Soc. Am. 54: 1535-1543 (1973).
- Young, E. and Sachs, M. B., Recovery of detection probability following sound exposure: comparison of physiology and psychophysics, J. Acoust. Soc. Am. 54: 1544-1553 (1973).
- Young, E. D., Recovery from Sound Exposure -- A Comparison of Psychophysics and Physiology, Ph.D. Thesis, Johns Hopkins University, 1972

2. Responses to Pure Tones

A. Rate versus level functions for auditory-nerve fibers for tone stimuli have been measured with a number of surprising new results:

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12 (7b).
Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

- i. There is a continuum of saturation behaviors related to fiber thresholds.
- ii. Slopes of rate-level functions depend on both the frequency of tone and threshold of the fibers.

These results are accounted for by a model relating basilar membrane displacement to fiber discharge rate via a saturating nonlinearity.

References:

Sachs, M. B. and Abbas, P. J., Rate versus level functions for auditory-nerve fibers in cats: tone-burst stimuli, J. Acoust. Soc. Am. 56: 1835-1847 (1974).

3. Responses to Tone-Tone Stimuli

An earlier stimulus-response relation for two-tone stimuli has been expanded. The cases considered were those where the ratio of component amplitudes were held constant. This case corresponds to the situation in a continuous spectrum signal (e.g., speech) whose spectrum is held constant while overall amplitude is varied.

Gross differences in behavior were found for second tones above and below fiber CF. These differences were well accounted for by an extension of our single-tone model.

References:

Abbas, P. J. and Sachs, M. B., Two-tone suppression in auditory-nerve fibers: extension of a stimulus-response relationship, J. Acoust. Soc. Am. 59: 112-122 (1976).

Abbas, P. J., Discharge Rate of Auditory-Nerve Fibers to Two-Tone Stimuli: An Experimental Study and a Model, Ph.D. thesis, Johns Hopkins University, 1974.

Sachs, M. B. and Abbas, P. J., Phenomenological model for two-tone suppression, J. Acoust. Soc. Am. 60: 1157-1163 (1976).

II. Coding of Stimuli in the Avian Cochlear Nerve

The first study of coding in the cochlear nerve was completed and published. This study showed that the stimulus transformations performed by the avian cochlea are similar in most respects to those performed by the mammalian cochlea.

Reference:

Sachs, M. B., Lewis, R. H., and Young, E. D., Discharge patterns of single fibers in the pigeon auditory nerve, Brain Research 70: 431-447 (1974).

III. Coding of Stimuli in the Avian Cochlear Nuclei

This project has been completed and is being prepared for publication. The major results are:

A. Nucleus magnocellularis presents a homogeneous population of cells, characterized by responses to tones similar to those of fibers in the cochlear nerve. The one exception is that the response areas of these cells are flanked by inhibitory sidebands. Tones in these sidebands inhibit the cells' spontaneous activity.

B. Cells in nucleus angularis present a more complex picture. There are at least three types based on response and inhibitory areas:

- i. Some cells show only excitatory responses.
- ii. Some cells show a central excitatory response area flanked by inhibitory sidebands.
- iii. Other cells show central inhibitory areas with complicated excitatory response bands.

There are at least four types of response patterns to tones in these cells. These are similar to those seen in mammalian cochlear nuclei:

- i. Primary-like
- ii. Pauser
- iii. Onset
- iv. Chopper

Reference:

Sachs, M. B. and Sinnott, J. M., Unit responses from the cochlear nuclei of the red-winged blackbird, Neuroscience Abstracts 2: 24 (1976).

Publications

Sachs, M. B., Quantitative studies of the responses of auditory-nerve fibers to two-tone stimuli, in Sachs, M. B. (Ed.), The Physiology of the Auditory System: A Workshop, National Educational Consultants, Baltimore, Maryland, pp. 113-124 (1971).

Young, E. and Sachs, M. B., Recovery from sound exposure in auditory-nerve fibers, J. Acoust. Soc. Am. 54: 1535-1543 (1973).

Young, E. and Sachs, M. B., Recovery of detection probability following sound exposure: comparison of physiology and psychophysics, J. Acoust. Soc. Am. 54: 1544-1553 (1973).

Sachs, M. B., Lewis, R. D., and Young, E. D., Discharge patterns of single fibers in the pigeon auditory nerve, Brain Research 7:- 431-447 (1974).

Sachs, M. B. and Abbas, P. J., Rate versus level functions for auditory-nerve fibers in cats: tone-burst stimuli, J. Acoust. Soc. Am. 56: 1835-1847 (1974).

Abbas, P. J. and Sachs, M. B., Two-tone suppression in auditory-nerve fibers: extension of a stimulus-response relationship, J. Acoust. Soc. Am. 59: 112-122 (1976).

Sachs, M. B. and Abbas, P. J., Phenomenological model for two-tone suppression, J. Acoust. Soc. Am. 60: 1157-1163 (1976).

Papers Presented

Young, E. D. and Sachs, M. B., Recovery of single auditory-nerve fibers from sound exposure, J. Acoust. Soc. Am. 50: 94A (1971).

Young, E. D. and Sachs, M. B., Recovery from sound exposure: comparison of human performance and responses from single auditory-nerve fibers in the cat, J. Acoust. Soc. Am. 52: 131A (1972).

Sachs, M. B., Lewis, R. H., and Young, E. D., Some properties of response patterns in the cochlear nerve of the pigeon, J. Acoust. Soc. Am. 52: 142A (1972).

Abbas, P. and Sachs, M. B., Two-tone suppression in cat auditory-nerve fibers: comparison of frequencies above and below fiber CF, J. Acoust. Soc. Am. 55: 466A (1974).

Sachs, M. B., Abbas, P. J. and Bevan, M. F., Discharge rate as a function of stimulus level for tones or noise in cat auditory-nerve fibers, J. Acoust. Soc. Am. 55: 466A (1974).

Sachs, M. B. and Sinnott, J. M., Unit responses from the cochlear nuclei of the red-winged blackbird, Neurosciences Abstracts 2: 24 (1976).

Papers Presented (continued)

Sinnott, J. M., Sachs, M. B. and Hienz, R. D., Differential sensitivity to frequency and intensity in songbirds, J. Acoust. Soc. Am. 60: S87 (1976).

Book

Sachs, M. B. (Ed.), The Physiology of the Auditory System: A Workshop, National Educational Consultants, Baltimore, Md. (1971).

Theses

Young, E. D., Recovery from Sound Exposure -- A Comparison of Psychophysics and Physiology, Ph.D. thesis, Johns Hopkins University, 1972.

Abbas, P. J., Discharge Rate of Auditory-Nerve Fibers to Two-Tone Stimuli: An Experimental Study and a Model, Ph.D. thesis, Johns Hopkins University, 1974